

Obstetrics & Gynecology

Accuracy of on-site tests to detect asymptomatic bacteriuria in pregnancy: a systematic review and meta-analysis

--Manuscript Draft--

Manuscript Number:	
Full Title:	Accuracy of on-site tests to detect asymptomatic bacteriuria in pregnancy: a systematic review and meta-analysis
Article Type:	Review
Manuscript Classifications:	Global health or global medicine; Prenatal care; Prenatal diagnosis
Corresponding Author:	Ewelina Rogozinska, MSc London, UNITED KINGDOM
Corresponding Author's Institution:	
First Author:	Ewelina Rogozinska, MSc
Order of Authors:	Ewelina Rogozinska, MSc Sandra Formina, MD Javier Zamora, Ph.D. Luciano Mignini, MD Khalid S. Khan, Msc
Manuscript Region of Origin:	UNITED KINGDOM

London, 22 April 2016

Dr. Nancy C. Chescheir
Editor-in-Chief
Obstetrics & Gynecology

Dear Dr. Chescheir

Re: “Accuracy of tests to detect asymptomatic bacteriuria during pregnancy: a systematic review and meta-analysis.”


Undetected and untreated bacteriuria during pregnancy is associated with poor pregnancy outcomes such as preterm delivery or low birth weight of the infant. Women with the undetected infection have a 20- 30-fold increased risk of developing pyelonephritis compared to women without the condition.

Although some bodies recommend a routine urine culture screening in early pregnancy, it is an expensive, cumbersome, and time-consuming test that requires access to laboratory facilities.

Our systematic review is the first comprehensive and robust synthesis of accuracy data concerning on-site tests to detect asymptomatic bacteriuria during antenatal care. We have identified publications from 27 primary studies (combined number of 13,641 women) with test accuracy data for nine tests suitable for use in the asymptomatic population. More than half of the studies came from low to upper-middle income countries (17/27).

Our work was commissioned by the World Health Organization to inform the new guidelines for antenatal care aiming to identify evidence-based interventions and procedures to improve pregnancy outcomes. We strongly feel that the work is relevant to the scope of Obstetrics & Gynecology, and we will be grateful if you could consider our manuscript for publication in your journal.

Yours sincerely



Ewelina Rogozińska on behalf of the manuscript authors


Women's Health Research Unit | Multidisciplinary Evidence Synthesis Hub (mEsh)
The Blizard Institute | Barts and the London School of Medicine and Dentistry
Queen Mary University of London, London

Friday, 22 April 2016

For the manuscript:

“Accuracy of on-site tests to detect asymptomatic bacteriuria in pregnancy: a systematic review and meta-analysis”

The lead author* affirms that this manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned (and, if relevant, registered) have been explained.

A handwritten signature in black ink, appearing to read 'Ewelina Rogozińska', with a large, stylized initial 'E'.

Signed by: Ewelina Rogozińska

*The manuscript's guarantor.

Accuracy of on-site tests to detect asymptomatic bacteriuria in pregnancy: a systematic review and meta-analysis

Ewelina Rogozińska MSc,^{1,2} Sandra Formina MD,³ Javier Zamora Ph.D.,^{1,2,4} Luciano Mignini MD,³ Khalid S. Khan MSc,^{1,2}

Affiliations

1. Women's Health Research Unit, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, London, UK
2. Multidisciplinary Evidence Synthesis Hub (mEsh), Barts and the London School of Medicine and Dentistry, Queen Mary University of London, London, UK
3. Centro Rosarino de Estudios Perinatales, Rosario, Argentina
4. Clinical Biostatistics Unit, Hospital Ramon y Cajal (IRYCIS) and CIBER Epidemiology and Public Health, Madrid, Spain

Corresponding author: Ewelina Rogozinska, Women's Health Research Unit, Centre for Primary Care and Public Health, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, tel: +44 20 7882 5881, e-mail: e.a.rogozinska@qmul.ac.uk

Funding The World Health Organization and the European Union funding to the EBM Connect Collaboration through Framework Programme, Marie Curie Actions, International Staff Exchange Scheme (Proposal no. 101377; Grant Agreement no. 247613).

Short Title: On-site tests for asymptomatic bacteriuria

Acknowledgments: The authors would like to acknowledge the assistance of the following advisors from the WHO Department of Reproductive Health and Research (A. Metin Gülmezoglu and Özge Tunçalp). We are thankful Dr. Caoimhe NicFhogartaigh, Microbiology Consultant at The Barts Health NHS Trust in London for her microbiological advice.

Conflict of interest: All authors have completed the Unified Competing Interest form an (available on request from the corresponding author) and declared that we have no conflicts of

26 interest. The views expressed are solely those of the authors and do not necessarily reflect the
27 decisions or stated policy of the World Health Organization.

28 **Precis:** Sensitivity of the on-site tests to detect asymptomatic bacteriuria in pregnancy
29 varies, however, their specificity to rule in the infection is high.

Abstract

Objective: The main objective of this systematic review of the literature was to determine the accuracy of on-site tests that require fewer resources to detect asymptomatic bacteriuria among pregnant women.

Data source: We searched the main electronic bibliographic databases from inception until June 2015 without language restrictions.

Methods of Study Selection: Two independent reviewers selected studies that recruited asymptomatic pregnant women to evaluate the accuracy of on-site tests in detecting the presence of bacteria in the urine using urine culture as a reference standard.

Tabulation, Integration, and Results: Data on women's characteristics, study design, sample collection, and handling were extracted along with to 2 x 2 tables, and synthesized, where possible, using a bivariate, hierarchical random effects model. Of 1,360 screened references, 27 papers (13,641 women) with test accuracy data on nine tests met the inclusion criteria. The most commonly evaluated test was urine dipstick. The pooled sensitivity and specificity of the dipstick to detect nitrites were 0.55 (95% CI 0.42 to 0.67) and 0.99 (95% CI 0.98 to 0.99), respectively. Griess test to detect nitrites had a sensitivity of 0.65 (95% CI 0.50 to 0.78) and specificity of 0.99 (95% CI 0.98 to 1.00). Dipslide with gram staining had a pooled sensitivity of 0.86 (95% CI 0.80 to 0.91) and specificity of 0.97 (95% CI 0.93 to 0.99).

Conclusions: The sensitivity of evaluated on-site tests to exclude bacterial urinary infection varies, however, their specificity to rule in disease is high.

Registration number: PROSPERO No. CRD42015027905

Keywords: test accuracy, asymptomatic bacteriuria, pregnancy, on-site test

Introduction

Asymptomatic bacteriuria, a common urinary tract infection, varies in prevalence by factors such as age, gender, level of sexual activity, etc. The prevalence of the infection in pregnancy ranges from 2–15% of whom 20–40% progress to symptomatic urinary infections (UTI).¹ Pregnant women with undetected asymptomatic bacteriuria are more likely to deliver prematurely² or low-birth-weight infants, and have a 20- 30-fold increased risk of developing pyelonephritis compared with those without the infection.³

Although some bodies recommend a routine urine culture screening in early pregnancy^{4, 5}, it is an expensive, cumbersome, and time-consuming test (taking 24 to 48 hours to obtain results) that requires access to laboratory facilities. There is a wide range of tests requiring fewer resources and minimal training,⁶ of which the most commonly used to detect the presence of bacteria instantly in the urine is a dipstick. Available evidence synthesis on their accuracy in pregnancy is limited in range of evaluated test⁷, and methodological strength.^{6, 8}

We bridge the above gap through a systematic and comprehensive evaluation of a wide range of on-site tests used to detect bacteriuria compared against urine culture as a reference standard in asymptomatic pregnant women taking into account potential sources of heterogeneity.

Methods

The review was conducted prospectively guided by a pre-defined protocol (PROSPERO No. CRD42015027905). We followed current standards of evidence synthesis for test accuracy⁹⁻¹¹ and reported findings in compliance with guidelines.¹²

Sources

We searched major databases such as Medline, Embase, Web of Science, Scopus, and a specialized database of Latin-American literature (LILACS) for studies published from database inception to August 2014, with no language restrictions. The search was updated to June 2015. The search strategy combined terms such as: ‘Pregnancy’, ‘Antenatal’, ‘Gestation’, ‘asymptomatic bacteriuria’ and ‘Urinary Tract Infections’ and applied a filter for test accuracy studies (for details see Appendix 1).¹³

Study selection

Two independent reviewers (ER and SF) screened references and full-text of previously selected articles. The consensus on the eligibility of evaluated publications was reached through discussion, or consultation with a third reviewer (KSK). We looked for studies reporting the accuracy of any on-site tests to detect asymptomatic bacteriuria among pregnant women without symptoms of urinary tract infections or not on antibiotic treatment. The reference standard had to be a urine culture, and asymptomatic bacteriuria had to be defined as equal, or more than 10⁵ Colony Forming Units of a single organism per mL of urine.⁸ Test accuracy had to be reported in a way allowing construction of 2 x 2 tables. We excluded studies with a case-control design and where reference standard was not reported or used a different definition of bacteriuria than specified above as this design and variation in reference standard were associated with bias.¹⁴

Data were extracted independently by ER and SF on to a piloted sheet. We collected authors’ details, year of publication, country, women’s characteristics, gestational age at testing; urine collection method, storage, and handling. The data were tabulated, cross checked and in the case of discrepancies discussed between the reviewers. The studies were grouped according to country income (low-, low-middle, upper-middle) using the World Bank classification.¹⁵ The

risk of bias and applicability of included studies were assessed by two independent reviewers (ER and SF) using the QUADAS-2 tool¹⁰ tailored for this review. Study quality was assessed for selection of participants, implementation of the index test and the reference standard, and patient flow. Studies with low risk of bias used a suitable spectrum of participants, recruited in consecutive or random manner; all participants were tested using the same reference standard, and the majority of the study population was included in analyses. Any disagreements over quality assessment were resolved by a third reviewer (KSK). We did not assess publication bias due to limitations of available methods.^{16, 17}

We calculated test accuracy estimates (sensitivity, specificity, and likelihood ratios for positive and negative test result) with 95% confidence intervals (CIs). Heterogeneity was investigated visually on forest plots with sensitivity and specificity estimates (with 95% confidence intervals) for individual studies. The impact of quality of study design, the reliability of population description, and sample collection and storage was explored through sensitivity analyses. All analyses were conducted using STATA version 12.1.¹⁸ If less than required number of data points was available, we pooled accuracy of sensitivity and specificity, and likelihood ratios using univariate model using *metaprop* and *metan* commands, respectively. Where a higher number of studies was available, we pooled the accuracy parameters using bivariate, random effects model as implemented in *metandi*¹⁹ and *midas*²⁰ commands. Posttest probabilities were calculated using following formula: $O = p1 / (1 - p1)$, $p2 = O * L$, $p = p2 / (1 + p2)$, where $p1$ pretest probability, O pretest odds, $p2$ posttest odds, L likelihood ratio, p posttest probability.²¹

Results

Out of the 1,360 references, 39 examining 27 types of index tests appeared initially to meet the inclusion criteria (Figure 1). After exclusion of tests not suitable for use in the asymptomatic population, we were left with 27 studies with nine index tests. List of all identified tests and reasons for study exclusion can be found in Appendix 2. Selected tests were: dipstick with only nitrites marker as positive, dipstick with nitrites or leucocytes as positive, urine analysis with bacteria count, dipslide with gram stain, Uricult (Orion Diagnostica, Espoo, Finland), Microstix-3 (Bayer Schering Pharma, Berlin, Germany), Griess test to detect nitrites, chlorhexidine reaction, and uriscreen catalase test. Reference of the included studies can be accessed in Appendix 3.

The majority of identified studies were conducted in low-middle (11 studies) or upper-middle (five studies) income countries; ten in high-income countries and only one in a low-income country. The studies were published between 1981 and 2015; ten studies were published before the year 2000, nine between 2000 and 2010 and remaining eight in the last five years. The majority (19/27) of included studies contributed to evidence synthesis accuracy data of only one test (Table 1) with urine dipstick as the most commonly reported test. Urine was mostly collected through clean catch midstream technique and as a random voided or first-morning sample in 56% of studies (15/27). Use of sterile containers was mentioned in ten out of 27 studies. More details on urine sample collection, handling and storage, and the details of urine culture incubation can be found in Appendix 4.

The overall quality of included studies was moderate (Figure 2). Twelve out of 27 studies gave a proper description of patients' selection with the remaining not giving enough details to assess this methodological aspect of the study. There was no concern for risk of bias due to index test implementation in over 80% of the studies (22/27). Similarly, for the reference

standard except two studies, the performance of the urine culture was classified as high risk of bias. Flow and timing were described with sufficient details in one-third of studies (9/27). The high concern over the applicability of findings was due to the type of the reference standard in five studies and the index test in one case. The main concern in the case of the reference standard was the use of a double urine culture to confirm the diagnosis of bacterial infection.

Twenty-one studies (9,491 women) reported accuracy data for the detection of nitrites using urine dipstick and eight for the combination of positive nitrites or leukocytes (5,940 women). The average prevalence of asymptomatic bacteriuria in these studies were 0.08 (95% CI 0.06 to 0.10). The pooled sensitivity of urine dipstick for positive nitrites in detecting infection was 0.55 (95% CI 0.42 to 0.67) with specificity 0.99 (95% CI 0.98 to 0.99). The pooled sensitivity of positive nitrites or leukocytes was 0.73 (95% CI 0.59 to 0.83) with specificity 0.89 (95% CI 0.79 to 0.94). For both tests, the accuracy parameters were heterogeneous with greater variability in sensitivity than specificity (Figure 3), 95% prediction contour was visibly wider for the combined markers (Appendix 5). The likelihood ratio of the positive test result for the urine dipstick test using only nitrites marker was 54.1 (95% CI 26.5 to 266.21).

One study each contributed data on the specificity and sensitivity of chlorhexidine reaction and uriscreen catalase tests. The sensitivity of the former was 1.00 (95% 0.65 to 1.00) and specificity (0.54, 95% CI 0.46 to 0.62) (Table 2). Use of Griess test to detect the presence of nitrites was reported in two studies (728 women). The sensitivity of the test was comparable to Uriscreen catalase test 0.65 (95% CI 0.50 to 0.78) with a specificity of 0.99 (95% CI 0.98 to 1.00). The likelihood of the positive test result was 56.6 (95% CI 12.6 to 255.1). Only one study reported the accuracy of the microscopic technique with the bacterial count in a centrifuged urine sample with a clearly defined threshold of more than 20 bacteria per High

Power Field (HPF). The sensitivity and specificity were 0.78 (95% CI 0.45 to 0.94) and 0.92 (95% CI 0.88 to 0.94), respectively.

Accuracy data of three dipslide-based tests included evaluation of Uricult (two studies), Microstix-3 (one study) and a generic dipslide method with gram stain dyeing and threshold of one or more bacteria per Oil Immersed Field (OIF) (six studies). Uricult had a sensitivity of 0.92 (95% CI 0.69 to 1.00) and specificity 0.85 (95% CI 0.24 to 1.00). The dipslide with gram staining on uncentrifuged urine had sensitivity and specificity of 0.86 (95% CI 0.80 to 0.91) and 0.97 (95% CI 0.93 to 0.99) respectively (Figure 3). The likelihood ratio of the positive test result was 30.2 (95% CI 11.9 to 76.6).

Sensitivity analysis was possible for dipstick with nitrites only as a marker, dipstick with nitrites or leukocytes and dipslide with gram staining. In all three cases, we explored the impact of population description and use of the sterile containers for urine storage. Neither of the factors changed the summary accuracy of the dipslide with gram staining. Analysis limited to studies with a clearly described population (asymptomatic women or not taking antibiotics) showed a marginal reduction in sensitivity (by 4%) for urine dipstick with positive leukocyte or nitrites marker. The pooled sensitivity of urine dipstick (nitrites with or without leukocytes) limited to studies providing details of urine container's sterility, presented a minimal increase in parameter precision. Findings from studies with low risk of bias and studies where the type of urine sample was not properly described had a minimal impact on the sensitivity the dipstick test with no change in the value of the pooled specificity.

Discussion

Out of 27 types of index tests identified in the literature, nine were suitable for use in the asymptomatic population. Three of them (urine dipstick, Griess test and dipslide with gram staining) had values of likelihood ratios for positive test result indicative of their usefulness (values > 10) in detecting asymptomatic bacteriuria during antenatal care. All test were minor to moderate usefulness to rule out the infection (likelihood ratios for negative result between 0.5–0.1).

This systematic review is the first comprehensive and robust synthesis of accuracy data concerning on-site tests to detect asymptomatic bacteriuria during antenatal care. Prospectively registered protocol with pre-specified population, reference standard, and definition of the outcome informed study selection, data extraction and analysis. On all stages of the review process, we followed current guidelines and standards.¹¹ The literature search in electronic databases restricted to test accuracy studies due to pragmatic reasons was supplemented by manual reference check. The publication bias due to limitations of available statistical methods^{16, 17} was not investigated in this review, however, we did undertake an extensive exploration of the heterogeneity between estimates of tests accuracy in individual studies.

The main limitation of this review was poor reporting in individual studies and paucity of data. The quality assessment was hindered by insufficient reporting of characteristics or recruited women, their flow through the study and timing between the use of index test and reference standard. Empirical evidence showed that test accuracy estimates can be affected by flaws in study design and its conduct.¹⁴ The estimates of test accuracy for four included tests were based on data from single studies with small sample sizes.²²⁻²⁴ This makes the parameters less reliable (wide confidence intervals) and more prone to chance findings. In order to compare the accuracy of all identified tests, we used the univariate model to pool sensitivity and specificity

estimates when less than four studies were available. Even though this approach does not account for correlation between two parameters as in the bivariate model, the findings should be fairly similar.²⁵ Despite these limitations our findings merit consideration as the most robust and current evidence synthesis.

The prevalence of asymptomatic bacteriuria in included studies ranged from 2 – 23% which overlaps with previously reported range¹. The likelihood ratios of the positive test result for the urine dipstick test (only nitrites), Griess test and generic dip slide with gram staining (bacterial count > 1/OIF) were indicative of tests usefulness in ruling in asymptomatic bacteriuria.²¹ The likelihood ratio of positive result with Dipslide Uricult due to wide confidence intervals cannot be considered reliable. However, its likelihood ratio for the negative result was the only one indicative its usefulness to rule out the infection (< 0.1). Likelihood ratios can be used to help adapt the results of the findings to individual situation basing on Bayes' theorem.²⁶ With pretest probability derived from identified studies we calculate the posttest probability of having the infection with a positive and negative test result (Table 2). Two out of nine evaluated tests (urine dipstick with positive nitrites and Griess test) increased the probability from 8.0% to above 80.0% in case of positive result, and both reduced it by half in case of a negative result. Need for training and access to basic laboratory facilities might make Griess test and Gram staining less attractive than urine dipstick in resource-limited settings.

Undetected and subsequently not treated asymptomatic bacteriuria is linked to pyelonephritis and other complications.³ Antibiotic treatment seem to reduce the risk of pyelonephritis in pregnancy and undesired pregnancy outcomes (preterm birth and low birth weight). Women incorrectly classified as positive (false positive) may be exposed to an unnecessary course of antibiotics with not well documented adverse effects.²⁷ In light of lack of robust evaluation of

harms and increasing antimicrobial resistance, it is crucial to correctly identify women who will truly benefit from the treatment.⁸

All identified on-site tests when positive increased posttest probability of detecting asymptomatic bacteriuria during the antenatal period. Urine dipstick, Griess test and dipslide with gram staining are most useful point-of-care options for ruling in the infection. Future research should aim to support the clinical decision-making on the management of asymptomatic pregnant women when access to urine culture is limited.

Contributors

ER selected eligible texts, data extraction form, extracted data, wrote the protocol, cleaned and analyzed the data, drafted and revised the manuscript. SF selected eligible texts, extracted data, and revised the paper. KSK, LM resolved discrepancies between reviewers and revised the draft paper. ER did statistical analysis, supervised by JZ. All authors contributed to the drafts and final version of the manuscript.

ER have full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Figures

Figure 1 Flow diagram describing selection of studies and tests

Figure 2 Study quality assessment using QUADAS-2 tool

Figure 3 Overview of sensitivity and specificity of tests to detect asymptomatic bacteriuria in pregnancy

277

278 **References**

- 279 1. Te West NIM, K.H. Urinary tract infection in gynaecology and obstetrics. *Obs Gyn Rep*
280 *Med* 2014; 24(11): 321-5.
- 281 2. Sheiner E, Mazor-Drey E, Levy A. Asymptomatic bacteriuria during pregnancy. *The*
282 *journal of maternal-fetal & neonatal medicine : the official journal of the European*
283 *Association of Perinatal Medicine, the Federation of Asia and Oceania Perinatal*
284 *Societies, the International Society of Perinatal Obstet* 2009; 22(5): 423-7.
- 285 3. Kincaid-Smith P, Bullen M. BACTERIURIA IN PREGNANCY. *Lancet (London,*
286 *England)* 1965; 1(7382): 395-9.
- 287 4. Council AHMA. Clinical Practice Guidelines: Antenatal Care – Module 1. Canberra,
288 Australia: Australian Government Department of Health and Ageing; 2012.
- 289 5. Excellence NifHaC. Antenatal care for uncomplicated pregnancies (CG62). In: Health
290 TNCCfWsaCs, editor. NICE guidelines: National Institute for Health and Care Excellence;
291 2008.
- 292 6. Lumbiganon P, Laopaiboon M, Thinkhamrop J. Screening and treating asymptomatic
293 bacteriuria in pregnancy. *Current opinion in obstetrics & gynecology* 2010; 22(2): 95-9.
- 294 7. Deville WL, Yzermans JC, van Duijn NP, Bezemer PD, van der Windt DA, Bouter LM.
295 The urine dipstick test useful to rule out infections. A meta-analysis of the accuracy. *BMC*
296 *urology* 2004; 4: 4.
- 297 8. Colgan R, Nicolle LE, McGlone A, Hooton TM. Asymptomatic bacteriuria in adults.
298 *American family physician* 2006; 74(6): 985-90.
- 299 9. Leeflang MM, Deeks JJ, Gatsonis C, Bossuyt PM. Systematic reviews of diagnostic test
300 accuracy. *Ann InternMed* 2008; 149(12): 889-97.

10. Whiting PF, Rutjes AW, Westwood ME, et al. QUADAS-2: a revised tool for the quality assessment of diagnostic accuracy studies. *Ann InternMed* 2011; 155(8): 529-36.
11. Deeks JJB, P.; Gatsonis, C. Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy. 2010.
12. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *Bmj* 2009; 339: b2700.
13. van der Weijden T, CJ IJ, Dinant GJ, van Duijn NP, de Vet R, Buntinx F. Identifying relevant diagnostic studies in MEDLINE. The diagnostic value of the erythrocyte sedimentation rate (ESR) and dipstick as an example. *Family practice* 1997; 14(3): 204-8.
14. Lijmer JG, Mol BW, Heisterkamp S, et al. Empirical evidence of design-related bias in studies of diagnostic tests. *Jama* 1999; 282(11): 1061-6.
15. Group TWB. Country and Lending Groups. 2015 2015.
<http://data.worldbank.org/about/country-and-lending-groups> (accessed 2/5/2016 2016).
16. Deeks JJ, Macaskill P, Irwig L. The performance of tests of publication bias and other sample size effects in systematic reviews of diagnostic test accuracy was assessed. *J Clin Epidemiol* 2005; 58(9): 882-93.
17. Song F, Khan KS, Dinnes J, Sutton AJ. Asymmetric funnel plots and publication bias in meta-analyses of diagnostic accuracy. *Int J Epidemiol* 2002; 31(1): 88-95.
18. StataCorp. Stata Statistical Software. 12.1 ed: College Station, TX: StataCorp LP. 2015.; 2015.
19. Harbord RM, Whiting P. metandi: Meta-analysis of diagnostic accuracy using hierarchical logistic regression. *The Stata Journal* 2009; 9(2): 211-29.
20. Dwamena BAS, R.; Carlos, R.C. Midas: meta-analysis of diagnostic accuracy studies. 2010.

21. Deeks JJ, Altman DG. Diagnostic tests 4: likelihood ratios. *BMJ (Clinical research ed)* 2004; 329(7458): 168-9.
22. Archbald FJ, Verma U, Tejani NA. Screening for asymptomatic bacteriuria with microstix. *Journal of Reproductive Medicine for the Obstetrician and Gynecologist* 1984; 29(4): 272-4.
23. Okusanya B, Aigere E, Eigbefoh J, Okome G, Gigi C. Is a chlorhexidine reaction test better than dipsticks to detect asymptomatic bacteriuria in pregnancy? *Journal of Obstetrics and Gynaecology* 2014; 34(1): 21-4.
24. Teppa RJ, Roberts JM. The uriscreen test to detect significant asymptomatic bacteriuria during pregnancy. *Journal of the Society for Gynecologic Investigation* 2005; 12(1): 50-3.
25. Simel DL, Bossuyt PM. Differences between univariate and bivariate models for summarizing diagnostic accuracy may not be large. *J Clin Epidemiol* 2009; 62(12): 1292-300.
26. Fagan TJ. Letter: Nomogram for Bayes theorem. *The New England journal of medicine* 1975; 293(5): 257.
27. Smaill FM, Vazquez JC. Antibiotics for asymptomatic bacteriuria in pregnancy. *The Cochrane database of systematic reviews* 2015; 8: Cd000490.

- 1 Table 1 Characteristics of studies evaluating accuracy of on-site tests to detect asymptomatic
- 2 bacteriuria in antenatal care settings

Study ID	Publication Year	Country	Number of women	Prevalence of the infection (95% CI)	Index test(s)
Abbasi 1985 ¹	1985	USA	65	0.05 (0.02–0.13)	Dipstick
Anandkumar 2011 ²	2011	India	300	0.13 (0.09–0.17)	Dipslide with gram staining
Archbald 1984 ³	1984	USA	287	0.03 (0.02–0.06)	Dipstick, Urinalysis (bacteria count), Dipslide (Microstix–3)
Awonuga 2011 ⁴	2011	Nigeria	205	0.11 (0.07–0.16)	Dipstick
Bachman 1993 ⁵	1993	USA	1047	0.02 (0.02–0.03)	Dipstick, Dipslide with gram staining
Balamurugan 2012 ⁶	2012	India	100	0.13 (0.08–0.21)	Dipstick
Campos-Outcalt 1985 ⁷	1993	USA	299	0.05 (0.03–0.08)	Dipstick
Demilie 2014 ⁸	2014	Ethiopia	330	0.08 (0.06–0.12)	Dipstick
Eigbefoh 2008 ⁹	2008	Nigeria	400	0.22 (0.18–0.26)	Dipstick
Gayathree 2010 ¹⁰	2010	India	900	0.07 (0.05–0.09)	Dipstick, Dipslide with gram staining
Graninger 1992 ¹¹	1992	Germany	1000	0.13 (0.11–0.15)	Dipstick
Greeff 2002 ¹²	2002	South Africa	247	0.23 (0.19–0.29)	Dipslide (Uricult)
Jayalakshmi 2008 ¹³	2008	India	630	0.07 (0.06–0.10)	Dipstick, Dipslide with gram staining

Kacmaz 2006 ¹⁴	2006	Turkey	250	0.04 (0.02–0.07)	Dipstick
Khattak 2004 ¹⁵	2004	Pakistan	290	0.06 (0.04–0.10)	Griess test
Kovavisarach 2008 ¹⁶	2008	Thailand	360	0.10 (0.07–0.14)	Dipstick
Mathews 1998 ¹⁷	1998	India	438	0.07 (0.05–0.10)	Griess test
Mignini 2009 ¹⁸	2009	Argentina	2353	0.14 (0.13–0.16)	Dipstick, Dipslide (Uricult)
Mukherjee 2015 ¹⁹	2015	India	250	0.08 (0.06–0.12)	Dipstick, Dipslide with gram staining
Okusanya 2014 ²⁰	2014	Nigeria	150	0.05 (0.02–0.09)	Dipstick, Chlorhexidine reaction
Pallarés 1990 ²¹	1990	Spain	74	0.12 (0.07–0.22)	Dipstick, Dipslide with gram staining
Plauche 1981 ²²	1981	USA	561	0.13 (0.09–0.18)	Dipstick
Shelton 2001 ²³	2001	USA	200	0.10 (0.07–0.15)	Dipstick
Soisson 1985 ²⁴	1985	USA	1062	0.06 (0.05–0.07)	Dipstick
Teppa 2005 ²⁵	2005	Venezuela	150	0.19 (0.13–0.26)	Uriscreen catalase test
Tincello 1998 ²⁶	1998	UK	893	0.05 (0.04–0.07)	Dipstick
Titoria 2014 ²⁷	2014	India	800	0.05 (0.04–0.07)	Dipstick

3 References of the included studies are available in *Appendix*

Table 1 Likelihood ratios and post-test probabilities for tests—detect asymptomatic bacteriuria in antenatal care settings

Index test	Likelihood ratio for a positive result (95% CI)	Probability* of infection after positive testing (%)	Likelihood ratio for a negative result (95% CI)	Probability* of infection after negative testing (%)
Dipstick: Nitrites positive	54.08 (26.50–266.21)	82.5	0.46 (0.35–0.61)	3.9
Dipstick: Leucocytes or nitrites positive	6.36 (3.31–12.21)	35.6	0.31 (0.19–0.49)	2.6
Chlorhexidine reaction	2.03 (1.58–2.61)	15.0	0.12 (0.01–1.71)	1.0
Uriscreeen catalase test	5.69 (3.15–10.32)	33.1	0.44 (0.27–0.70)	3.7
Griess test (nitrites)	56.62 (12.57– 255.06)	83.1	0.36 (0.25–0.53)	3.0
Urinalysis (bacteria count >20/HPF)	9.4 (5.56–15.89)	44.9	0.24 (0.07–0.82)	2.1
Dipslide with gram staining (bacteria count ≥1/OIF)	30.22 (11.91–76.64)	72.4	0.14 (0.09–0.21)	1.2
Dipslide (Uricult)	20.04 (0.01–5.2e+04)	63.5	0.09 (0.00–2.16)	0.8
Dipslide (Microstix-3)	9.07 (5.21–15.77)	44.1	0.36 (0.18–0.74)	3.0

*Pretest probability of 8.0% was a median prevalence derived from included studies

Figure 1 Study selection





Figure 1 Flow diagram describing studies and tests selection stages

Figure 2 Quality assessment



**For Okusanya 2014 studies risk of bias for Chlorhexidine test unclear*

Figure 2 Study quality assessment using QUADAS-2 tool. Number of studies classified as low (*light grey*), unclear (*dark grey*) or high (*black*) risk of bias/concern for applicability

Figure 3 Overview of tests accuracy




Figure 3 Overview of sensitivity and specificity of included tests to detect asymptomatic bacteriuria among pregnant women (for details see *Appendix 4*)

Appendix 1 Search strategy in MEDLINE via Ovid (19th Aug 2014 repeated on 2nd June 2015

Item Search term

1. Pregnancy.mp. or exp Pregnancy/
2. exp Gravidity/
3. gravid*.mp.
4. gestation*.mp.
5. exp Pregnant Women/
6. pregnant wom#n.mp.
7. (child adj3 bearing).mp.
8. childbearing.mp.
9. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8
10. exp sensitivity/
11. exp specificity/
12. 10 and 11
13. (predictive adj3 value\$).tw.
14. (sensitivity or specificity).tw.
15. exp Diagnostic Errors/
16. ((false adj positiv\$) or (false adj negativ\$)).tw.
17. (observer adj variation\$).tw.
18. (roc adj curve).tw.
19. (likelihood adj3 ratio\$).tw.
20. exp Likelihood Functions/
21. 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20
22. exp bacteriuria/
23. (asymptomatic\$ adj2 bacteriuria\$).tw.
24. exp Urinary Tract Infections/
25. 22 or 23 or 24
26. 9 and 21 and 25

Appendix 2 Overview of test to detect asymptomatic bacteriuria reported in the retrieved literature

Group	No.	Index test	Decision	Justification for exclusion
Dipstick	1	Dipstick (marker: nitrites)	Include	
	2	Dipstick (marker: leucocytes or nitrites)	Include	
	3	Dipstick (marker: leucocytes)	Exclude	Population: Information not useful in asymptomatic population
	4	Dipstick (marker: leucocytes and nitrites)	Exclude	Population: Information not useful in asymptomatic population
Culture methods (Dip slides)	5	Uricult & Uricult Trio (Orion Diagnostica)^	Include	
	6	Testuria (Ayerst Labs)	Exclude	Relevance to clinical practice: irrelevance to current clinical practice
	7	Bacturcult (Wampole Labs)	Exclude	Relevance to clinical practice: irrelevance to current clinical practice
	8	Microstix-3	Include	
	9	Generic dipslide	Exclude	Relevance to clinical practice: lack of information about threshold
Microscopic techniques	10	Microscopic analysis of urine (marker & threshold: >20 bacteria per High Power Field)	Include	
	11	Dip slide with gram staining (marker & threshold: ≥ 1 bacteria per oil immersed field)	Include	
	12	Microscopic analysis of urine (marker: leucocytes)	Exclude	Population: Not suitable to use in asymptomatic population

Group	No.	Index test	Decision	Justification for exclusion
Microscopic techniques	13	Dip slide with gram staining (threshold ≥ 2 bacteria per oil immersed field)	Exclude	Relevance to clinical practice: Excluded due to a threshold
Tests not usually used to detect bacteriuria	14	Uriscreen catalase tests (Savyon Diagnostics)	Include	
	15	Chlorhexidine reaction	Include	
Tests not usually used to detect bacteriuria	16	Uriglox (glucose level)	Exclude	Relevance to clinical practice: Test used to detect the presence or absence of the small physiological amount of glucose (2-20 mg per 100 ml)
	17	Griess test (test to detect nitrites)	Include	
	18	Lumac/3M Bacteriuria Screening Kit RLU>200	Exclude	Relevance to clinical practice: The test procedure is based upon firefly luciferase analysis of bacterial ATP. Exclude from further work as it requires lab facility
	19	Interleukin-8 ≥ 264 pg/mL	Exclude	Relevance to clinical practice: Interleukin-8 (IL-8), an inflammatory cytokine, is involved in host response to infection through neutrophil chemo attraction. Exclude from further work as it requires lab facility
	20	Chromogenic limulus assay	Exclude	Relevance to clinical practice: Not taken further due to identification of only gram negative pathogens
	21	Catalase tests	Exclude	Data concern: Subgroup analysis limits spectrum
	22	Triphenyl Tetrazolium Chloride	Exclude	Data concern: Subgroup analysis limits spectrum

Group	No.	Index test	Decision	Justification for exclusion
Automatic analysis	23	Automatic analysis of urine samples (marker & threshold: nitrites)	Exclude	Cost: Automated urine analysis is not widely available in resource limited settings
	24	Automatic analysis of urine samples (marker & threshold: >1200 bacteria per ml)	Exclude	Cost: Automated urine analysis is not widely available in resource limited settings
	25	Automatic analysis of urine samples (marker & threshold: leukocytes)	Exclude	Population: Information not useful in asymptomatic population
	26	Automatic analysis of urine samples (marker & threshold: leukocytes and nitrites)	Exclude	Population: Information not useful in asymptomatic population
	27	Automatic analysis of urine samples (Bac-T-Screen, Marion Labs)	Exclude	Relevance to clinical practice: Instrument is designed to provide a rapid, semi quantitative measurement of the bacteria present in urine specimens. Exclude from further work due to irrelevance to current clinical practice

Appendix 5 Reference list of all included studies

1. Abbasi IA, Hess LW, Johnson TR, McFadden E, Chernow B. Leukocyte esterase activity in the rapid detection of urinary tract and lower genital tract infections in obstetric patients. *American Journal of Perinatology*. 1985;2(4):311-3.
2. Anandkumar H, Srinivasa H, Kodliwadmah S, Raksha R. Symptomatic and Asymptomatic Urinary Tract Infection by *Escherichia coli* Among Pregnant Women Attending Out Patient Clinic of Obstetrics and Gynecology. *Journal of Pure and Applied Microbiology*. 2011;5(2):717-23.
3. Archbald FJ, Verma U, Tejani NA. Screening for asymptomatic bacteriuria with microstix. *Journal of Reproductive Medicine for the Obstetrician and Gynecologist*. 1984;29(4):272-4.
4. Awonuga DO, Fawole AO, Dada-Adegbola HO, Olola FA, Awonuga OM. Asymptomatic bacteriuria in pregnancy: evaluation of reagent strips in comparison to microbiological culture. *African Journal of Medicine & Medical Sciences*. 2011;40(4):377-83.
5. Bachman JW, Heise RH, Naessens JM, Timmerman MG. A study of various tests to detect asymptomatic urinary tract infections in an obstetric population. *Journal of the American Medical Association*. 1993;270(16):1971-4.
6. Balamurugan S, Shah C, Jayapriya S, Priyadarshini S, Jeya M, Ramesh RK. Reagent strip testing (RST) for asymptomatic bacteriuria (ASB) in pregnant women: A cost-effective screening tool in under-resourced settings. *Journal of Clinical and Diagnostic Research*. 2012;6(4 SUPPL. 2):671-3.
7. Campos-Outcalt DE, Corta PJ. Screening for asymptomatic bacteriuria in pregnancy. *The Journal of family practice*. 1985;20(6):589-91.
8. Demilie T, Beyene G, Melaku S, Tsegaye W. Diagnostic accuracy of rapid urine dipstick test to predict urinary tract infection among pregnant women in Felege Hiwot Referral Hospital, Bahir Dar, North West Ethiopia. *BMC research notes*. 2014;7:481.
9. Eigbefoh J, Isabu P, Okpere E, Abebe J. The diagnostic accuracy of the rapid dipstick test to predict asymptomatic urinary tract infection of pregnancy. *Journal of Obstetrics and Gynaecology*. 2008;28(5):490-5.
10. Gayathree L, Shetty S, Deshpande SR, Venkatesha DT. Screening for asymptomatic bacteriuria in pregnancy: An evaluation of various screening tests at the hassan district hospital, India. *Journal of Clinical and Diagnostic Research*. 2010;4(4):2702-6.
11. Graninger W, Fleischmann D, Schneeweiss B, Aram L, Stockenhuber F. Rapid screening for bacteriuria in pregnancy. *Infection*. 1992;20(1):9-11.
12. Greeff A, Jeffery B, Pattinson RC. Uricult trio as a screening test for bacteriuria in pregnancy. *South African Medical JournalSuid-Afrikaanse Tydskrif Vir Geneeskunde*. 2002;92(4):306-9.
13. Jayalakshmi J, Jayaram V. Evaluation of various screening tests to detect asymptomatic bacteriuria in pregnant women. *Indian Journal of Pathology and Microbiology*. 2008;51(3):379-81.
14. Kacmaz B, Cakir O, Aksoy A, Biri A. Evaluation of rapid urine screening tests to detect asymptomatic bacteriuria in pregnancy. *Japanese Journal of Infectious Diseases*. 2006;59(4):261-3.
15. Khattak AMK, H.; Akhtar, W.;Mahsud, I.;Ashiq, B. Accuracy of griess test to predict asymptomatic bacteriuria during pregnancy. *Gomal Journal of Medical Sciences*. 2004;2(1):20-3.
16. Kovavisarach E, Vichaipruck M, Kanjanahareutai S. Reagent strip testing for antenatal screening and first meaningful of asymptomatic bacteriuria in pregnant women. *Journal of the Medical Association of Thailand*. 2008;91(12):1786-90.
17. Mathews JE, George S, Mathews P, Mathai E, Brahmadathan KN, Seshadri L. The Griess test: An inexpensive screening test for asymptomatic bacteriuria in pregnancy. *Australian & New Zealand Journal of Obstetrics & Gynaecology*. 1998;38(4):407-10.

18. Mignini L, Carroli G, Abalos E, Widmer M, Amigot S, Nardin JM, et al. Accuracy of diagnostic tests to detect asymptomatic bacteriuria during pregnancy. *Obstetrics & Gynecology*. 2009;113(2:Pt 1):t-52.
19. Mukherjee KG, S.; Babita, V.C.L.; Bhattacharjee, D.; Chakroborti, G. A study on asymptomatic bacteriuria in pregnancy: prevalence, etiology and comparison of screening methods. *Int J Res Med Sci*. 2014;2(3):1085-91.
20. Okusanya B, Aigere E, Eigbefoh J, Okome G, Gigi C. Is a chlorhexidine reaction test better than dipsticks to detect asymptomatic bacteriuria in pregnancy? *Journal of Obstetrics and Gynaecology*. 2014;34(1):21-4.
21. Pallares J, Casas J, Guarga A, Marquet R, Solans P, Muxi C, et al. The evaluation of different methods for rapid diagnosis in the detection of asymptomatic bacteriuria in pregnant women. *Atencion primaria / Sociedad Espanola de Medicina de Familia y Comunitaria*. 1990;7(9):547-50.
22. Plauche WC, Janney FA, Curole DN. Screening for asymptomatic bacteriuria in pregnant patients: three office screening systems versus quantitative culture. *Southern Medical Journal*. 1981;74(10):1227-9.
23. Shelton SD, Boggess KA, Kirvan K, Sedor F, Herbert WNP. Urinary interleukin-8 with asymptomatic bacteriuria in pregnancy. *Obstetrics and Gynecology*. 2001;97(4):583-6.
24. Soisson AP, Watson WJ, Benson WL, Read JA. Value of a screening urinalysis in pregnancy. *Journal of Reproductive Medicine*. 1985;30(8):588-90.
25. Teppa RJ, Roberts JM. The uriscreen test to detect significant asymptomatic bacteriuria during pregnancy. *Journal of the Society for Gynecologic Investigation*. 2005;12(1):50-3.
26. Tincello DG, Richmond DH. Evaluation of reagent strips in detecting asymptomatic bacteriuria in early pregnancy: prospective case series. *BMJ*. 1998;316(7129):435-7.
27. Titoria A, Gupta A, Rathore AM, Prakash SK, Rawat D, Manaktala U. Asymptomatic bacteriuria in women attending an antenatal clinic at a tertiary care centre. *South African Journal of Obstetrics and Gynaecology*. 2014;20(1):4-7.

Appendix 4 Details of urine sample collection, processing and storage in individual studies

No.	Study ID	Description of the population	Urine collection	Random voided or first morning sample	Description of urine container	Time before sending to lab	Sample storage	Sample centrifuging	Max. time of sample storage	T° Incubation of urine culture	Hours incubation of urine culture
1	Abbasi 1985	women not on antibiotics	clean catch midstream specimens	unspecified	unspecified	unspecified	unspecified	centrifuged	unspecified	unspecified	unspecified
2	Anandkumar 2011	asymptomatic women	clean catch midstream specimens	unspecified	sterile	1 hour	unspecified	uncentrifuged	unspecified	37°C	24 hours
3	Archbald 1984	lack of detailed information	clean catch midstream specimens	random voided, at place of care	unspecified	3 hours (for IT at place), samples sent to microbiology lab for UC	unspecified	centrifuged	unspecified	unspecified	unspecified
4	Awonuga 2011	asymptomatic women	clean catch midstream specimens	unspecified	sterile	immediately	4°C	unspecified	unspecified	unspecified	unspecified
5	Bachman 1993	women not on antibiotics	clean catch midstream specimens	random voided, at place of care	unspecified	45 minutes	unspecified	Dipstick: unspecified; Slide with gram staining: uncentrifuged	unspecified	unspecified	unspecified
6	Balamurugan 2012	asymptomatic women	clean catch midstream specimens	random voided, at place of care	sterile	1 hour	refrigerated	centrifuged	4 hours	unspecified	unspecified
7	Campos Outcalt 1985	lack of detailed information	midstream specimens	random voided, at place of care	sterile	unspecified	unspecified	unspecified	unspecified	unspecified	unspecified
8	Demilie 2014	asymptomatic women	clean catch midstream specimens	random voided, at place of care	sterile	2 hours	4°C	unspecified	unspecified	37°C aerobically	24 hours

No.	Study ID	Description of the population	Urine collection	Random voided or first morning sample	Description of urine container	Time before sending to lab	Sample storage	Sample centrifuging	Max. time of sample storage	T° Incubation of urine culture	Hours incubation of urine culture
9	Eigbefoh 2008	asymptomatic women	clean catch midstream specimens; catheterization (8 women)	random voided, at place of care	sterile	1 hour	unspecified	uncentrifuged	unspecified	37°C	24 hours
10	Gayathree 2010	lack of detailed information	clean catch midstream specimens	unspecified	sterile	1 hour	refrigerated	uncentrifuged	unspecified	35°C aerobically	18-24 hours
11	Graninger 1992	women not on antibiotics	clean catch midstream specimens	unspecified	unspecified	30 minutes	unspecified	unspecified	unspecified	35°C	18 hours
12	Greeff 2002	asymptomatic women	midstream specimens	random voided, at place of care	unspecified	unspecified	unspecified	uncentrifuged	unspecified	unspecified	unspecified
13	Jayalakshmi 2008	lack of detailed information	clean catch midstream specimens	unspecified	unspecified	1 hour	unspecified	uncentrifuged	unspecified	37°C aerobically	24 hours
14	Kacmaz 2006	asymptomatic women	clean catch midstream specimens	unspecified	sterile	1 hour	4°C	unspecified	4 hours	unspecified	12-24-48 hours
15	Khattak 2004	lack of detailed information	clean catch midstream specimens	random voided, at place of care	sterile	immediately	unspecified	unspecified	unspecified	37°C	unspecified
16	Kovavisarath 2008	asymptomatic women	clean catch midstream specimens	random voided, at place of care	sterile	30 minutes	unspecified	unspecified	unspecified	35-37°C	24-48 hours
17	Mathews 1998	lack of detailed information	clean catch midstream specimens	random voided, at place of care	unspecified	immediately	unspecified	unspecified	unspecified	unspecified	unspecified

No.	Study ID	Description of the population	Urine collection	Random voided or first morning sample	Description of urine container	Time before sending to lab	Sample storage	Sample centrifuging	Max. time of sample storage	T° Incubation of urine culture	Hours incubation of urine culture
18	Mignini 2009	asymptomatic women	clean catch midstream specimens	random voided, at place of care	sterile	1 hour	4°C	unspecified	4 hours	37°C	24 hours
19	Mukherjee 2014	asymptomatic women	clean catch midstream specimens	unspecified	sterile	1 hour	4°C	uncentrifuged	24 hours	37°C aerobically	24-48 hours
20	Okusanya 2014	asymptomatic women	clean catch midstream specimens	unspecified	sterile	1 hour	4°C	Dipstick: uncentrifuged; Chlorhexidine reaction: unspecified	unspecified	unspecified	unspecified
21	Pallarés 1990	women not on antibiotics	clean catch midstream specimens	first morning	sterile	90 minutes	4°C	unspecified	unspecified	35-35°C	24 hours
22	Plauche 1981	asymptomatic women	clean catch midstream specimens	unspecified	sterile	30 minutes	unspecified	unspecified	unspecified	unspecified	unspecified
23	Shelton 2001	lack of detailed information	clean catch midstream specimens	unspecified	unspecified	immediately	unspecified	unspecified	unspecified	37°C	24 hours
24	Soisson 1985	lack of detailed information	clean catch midstream specimens	unspecified	unspecified	unspecified	unspecified	unspecified	unspecified	unspecified	unspecified
25	Teppa 2005	asymptomatic women	in-and-out sterile technique of catheterization	first morning or random voided 4 hours of incubation	sterile	unspecified	4°C	unspecified	4 hours	unspecified	unspecified

No.	Study ID	Description of the population	Urine collection	Random voided or first morning sample	Description of urine container	Time before sending to lab	Sample storage	Sample centrifuging	Max. time of sample storage	T° Incubation of urine culture	Hours incubation of urine culture
26	Tincello 1998	asymptomatic women	midstream specimens	random voided, at place of care	unspecified	unspecified	unspecified	unspecified	unspecified	unspecified	unspecified
27	Titoria 2014	asymptomatic women	clean catch midstream specimens	random voided	sterile	4 hours	unspecified	uncentrifuged	unspecified	unspecified	unspecified

Appendix 5 Overview of sensitivity and specificity reported in studies included in the review by test




Figure 1 Overview of sensitivity and specificity in individual studies reporting use of dipstick test (only nitrites marker positive) in detecting asymptomatic bacteriuria among pregnant women




Figure 2 Receiver operating characteristic plots for studies evaluating accuracy of dipstick test (only nitrites marker positive) in detecting asymptomatic bacteriuria among pregnant women




Figure 3 Overview of sensitivity and specificity in individual studies reporting use of dipstick test (nitrites or leucocytes marker positive) in detecting asymptomatic bacteriuria among pregnant women




Figure 4 Receiver operating characteristic plots for studies evaluating accuracy of dipstick test (nitrites or leucocytes marker positive) in detecting asymptomatic bacteriuria among pregnant women






Figure 5 Overview of sensitivity and specificity in individual studies reporting use of dipslide with gram staining (more than one bacteria per OIF) in detecting asymptomatic bacteriuria among pregnant women


Chlorhexidine reaction




Uriscreeen catalase test




Griess test



Urinalysis: bacteria count >20/HPF



Dipslide: Uricult



Dipslide: Microstix-3




Figure 6 Overview of sensitivity and specificity in individual studies reporting use of remaining tests in detecting asymptomatic bacteriuria among pregnant women

Author Agreement

Corresponding author:

Ewelina Rogozinska Women's Health Research Unit

Name

Address

+44 207 882 5831

e.a.rogozinska@gmail.ac.uk

Telephone number

FAX number

E-mail address

**Title of Manuscript: Accuracy of on-site test to detect asymptomatic bacteriuria

**Name(s) of author(s): In pregnancy: a systematic review and meta-analysis

Rogozinska, Formica, Zomara, Mignini, Khan.

A signature below certifies agreement with the corresponding author designation and acceptance of this agreement. If there is more than one author, each must sign this agreement.

I. CONDITIONS OF SUBMISSION

A. Scope of Agreement: Unless otherwise specified, the terms of this Agreement apply to the Manuscript and any and all supplemental digital content (SDC) (audio, video, and other data files, the forms of which may vary) submitted by the author(s) for online publication with the Manuscript. The Manuscript and the SDC hereinafter are referred to collectively as the "Work."

B. Sole Submission of Original Work: Each author represents and warrants that the Work is the author's original work (except for any material from previously copyrighted works obtained by permission or assignment and credited as such); that none of the information has been published previously or is under consideration by another publication; that upon publication, the Work shall not infringe any copyright, invade any right of privacy, contain any libelous material, or infringe or violate any other right of any person or entity. Each author represents and warrants that the following is a complete description of all material contained in the Work, including any SDC, which is not original and/or in which the authors are not able to transfer copyright (attach additional pages if necessary):

C. Approval Verification: Each author represents and warrants that investigations involving human or animal subjects received the appropriate institutional approval and agrees to provide evidence of such approval if requested. In addition, each author agrees to produce the data on which the Work is based for examination by the editors or their assignees should they request them.

D. Authorship Certification: Each author represents and warrants that he or she has fulfilled the definition of authorship given in the Instructions for Authors, participated in this study sufficiently to ensure its integrity and to take public responsibility for it, and approved the Work submitted to *Obstetrics & Gynecology*. Each author represents and warrants that all persons who contributed to the work reported in the Manuscript are identified as authors or are included in the acknowledgements.

E. Conflict of Interest Disclosure: Each author represents and warrants that he or she has no financial affiliation (eg, employment, direct payments, stock holdings, retainers, consultantships, patent-licensing arrangements, or honoraria) or involvement within the last 3 years with any commercial organization with potential financial interest in the subject or materials discussed in the Work, except as disclosed in an attachment to this document. Any other potential conflict of interest also is disclosed.

F. Journal Research: Works under consideration are sometimes enrolled in *Obstetrics & Gynecology* research projects. When this is done, the identity of individual Works and authors is kept confidential. Each author hereby give consent for the Work to be used in such research.

G. Compliance with Privacy Laws: Each author represents and warrants that the Work complies with all applicable patient privacy laws (such as the Health Insurance Portability and Accountability Act of 1996, the regulations promulgated thereunder, and the Health Information Technology for Economic and Clinical Health Act of 2009, if applicable) and that the authors have obtained, prior to submission, any necessary patient authorizations. The corresponding author shall maintain such authorizations and shall inform the journal in a timely manner of any revocation by a patient of such authorization.

H. Identification of Patients: Each author represents and warrants that he or she has obtained a signed consent form from each patient described in a case report and each patient who is identifiable in a photograph or video. *Obstetrics & Gynecology* reserves the right to review the signed forms. Should the editorial office request copies of the signed forms, the corresponding author shall submit the forms in a timely manner.

I. Acknowledgments: Each author represents and warrants that all persons who contributed to the work reported in the Manuscript, but not sufficiently to be authors, are named in the acknowledgments and have given written permission to be named.

J. Manuscript Preparation Assistance: Each author represents and warrants that any and all document preparation assistance, including but not limited to topic development, data collection, analysis, writing, or editorial assistance, is disclosed in the acknowledgments.

K. Consideration: Each author hereby acknowledges that the publication of the Manuscript in a forthcoming edition of *Obstetrics & Gynecology* constitutes the sole consideration due to each of the authors under this Agreement.

OBSTETRICS & GYNECOLOGY

I. Indemnification and Hold Harmless: Each author hereby agrees to indemnify, defend, and hold harmless the American College of Obstetricians and Gynecologists (the College), its directors, officers, agents, employees, affiliates, successors and assigns from and against all claims, actions, suits or other proceedings, and any and all losses, judgments, settlements, damages, expenses, liabilities (including liability assumed in contractual agreements with third parties), or other costs (including reasonable counsel fees, disbursements, and costs of settlement), arising from or in any way relating to the Work, or any breach, or alleged breach by any author of any of the provisions of this Agreement. If the Authors fail to promptly or diligently pursue any defense of any indemnified party, the indemnified parties, or any of them, may assume such defense at the expense of the Authors.

II. COPYRIGHT

A. Copyright Transfer: Each author hereby assigns, transfers, and otherwise conveys all rights, title, and interest in the Work, including all copyright ownership worldwide, in all languages, and in all forms of media now or hereafter known, including electronic media such as CD-ROM, Internet, and Intranet, for the full term and all extensions and renewals thereof, to the College. Each author acknowledges that, in addition to publishing the Work in *Obstetrics & Gynecology*, the College may use, publish, and distribute the Work in any manner or media in its sole discretion including, but not limited to, licensing the SDC to YouTube for inclusion on *Obstetrics & Gynecology's* YouTube channel. If the College and the editor should decide for any reason not to publish the Work in *Obstetrics & Gynecology*, one of them shall give prompt notice of the decision to the corresponding author, and this Agreement and each of the rights and obligations described herein shall terminate.

B. Work Made For Hire: If this Work, or any portion of it, was written in the course of the author(s)' employment as a "Work Made for Hire" as defined under U.S. Copyright Law, the Work is owned by the company/employer. The author must check the "Work for Hire" box after his or her signature and an authorized representative of the company/employer must also sign this form and state his or her title. In such case, the company/employer hereby assigns to the College, during the full term of copyright, all copyright in and to the submitted Work for the full term of copyright throughout the world as specified in Section II, paragraph A above.

C. Government Employees: If any author created this Work, or a portion of it, in the course of that author's employment by the United States Government, that author must check the "Government" box after his or her signature. A work prepared by a government employee as part of his or her official duties is called a "work of the U.S. Government" as it is not subject to copyright and the provisions of this Section II, paragraphs A and B, do not apply. If it is not prepared as part of the employee's official duties, it may be subject to copyright.

III. DEPOSIT OF POSTPRINTS

A. Author(s) Posting of Articles to an Institutional Repository: *Obstetrics & Gynecology* will permit the author(s) to deposit a "postprint" (defined as the final Manuscript after peer-review and acceptance for publication but prior to the publisher's copyediting, design, formatting, and other services) on his/her personal Web site, university's institutional repository, or employer's intranet, subject to the following:

1. The author(s) may only deposit the postprint, as designated by *Obstetrics & Gynecology*, and not any other version of the Manuscript.
2. The author(s) shall not update the postprint text or replace it with a proof or with the final published version.
3. Except as provided in this Section, the author(s) shall not include the postprint or any other version of the article in any commercial website or in any repository other than as specified above.
4. The author(s) shall not display the postprint until 12 months after publication of the final article.
5. The author(s) must attach the following notice to the postprint: "This is a non-final version of an article published in final form in (provide complete journal citation)."
6. The author(s) shall provide a link in the postprint to the *Obstetrics & Gynecology* Internet site.

B. Compliance With Certain Research Funding Agency Requirements:

1. Authors of articles based on research funded by the National Institutes of Health (NIH), the Howard Hughes Medical Institute (HHMI), or the Wellcome Trust, are subject to the following policies:
 - a. NIH requires authors to deposit postprints based on NIH-funded research in the National Library of Medicine's (NLM) PubMed Central (PMC) within 12 months after publication of the final article.
 - b. HHMI requires authors to deposit postprints based on HHMI-funded research in NLM's PMC within 6 months after publication of the final article.
 - c. The Wellcome Trust requires authors to deposit postprints based on research funded by the Wellcome Trust in UK PubMed Central within 6 months after publication of the final article.
2. If the author(s) indicate at the bottom of this Agreement that the Manuscript is subject to a public access requirement, LWW will transmit the postprint of the Manuscript to NLM's PMC. The author(s) agree to the following:
 - a. Upon NIH request, it remains the legal responsibility of the author(s) to confirm with NIH the provenance of the Manuscript for purposes of deposit.
 - b. The author(s) shall not deposit their articles themselves.
 - c. The author(s) shall not alter the postprint already transmitted to NLM's PMC.

C. "Public Access Policy" Funding Disclosure

Please disclose below if you have received funding for research on which your article is based from any of the following organizations

- ☐ National Institutes of Health (NIH)
- ☐ Wellcome Trust
- ☐ Howard Hughes Medical Institute (HHMI)
- ☐ Other (please list):
- ☐ Other (please list):

Each author must complete and sign a copy of this form to indicate his/her acceptance of the terms of this Agreement. Please check the applicable boxes.

Author:

Check if applicable (see Section II)

- ☐ Work for Hire (see Section II.B). Representative of company/employer must sign below.

Company/employer representative:	<input type="text"/>	<input type="text"/>	<input type="text"/>
	Signature	Printed Name	Date
	<input type="text"/>	<input type="text"/>	
	Title	Company/Employer Name	

- ☐ Government (see Section II.C)

Disclosure of Potential Conflicts of Interest

- ☐ Yes, I have disclosed potential conflicts of interest (attached) (see Section I.E)
- ☒ No, I have no potential conflicts of interest to disclose (see Section I.E)

Authorship

In January 2014, *Obstetrics & Gynecology* adopted the ICMJE recommendations for authorship. *Note that you must meet a minimum of four criteria to be an author.* Individuals who do not qualify for authorship based on the ICMJE recommendations will be listed in the Acknowledgments. Indicate your contributions to the Work by checking the applicable boxes:

- ☐ 1a) Substantial contributions to the conception or design of the Work
- or
- ☐ 1b) Substantial contributions to the acquisition, analysis, or interpretation of data for the Work
- and
- ☐ 2) Drafting the Work or revising it critically for important intellectual content
- ☒ 3) Final approval of the version to be published
- ☐ 4) Agreement to be accountable for all aspects of the Work in ensuring that questions related to the accuracy or integrity of any part of the Work are appropriately investigated and resolved

Author Signature

Printed Name

Date

MIGNINI LUCIANO

26.04.16

Author Agreement

Corresponding author:

Ewelina Rogozinska

Women's Health Research Unit

Name

Address

+44 207 882 5831

e.a.rogozinska@gmail.ac.uk

Telephone number

FAX number

E-mail address

**Title of Manuscript: Accuracy of on-site test to detect asymptomatic bacteriuria

**Name(s) of author(s): in pregnancy: a systematic review and meta-analysis
Rogozinska; Kormia; Zomara; Mignini; Khan.

A signature below certifies agreement with the corresponding author designation and acceptance of this agreement. If there is more than one author, each must sign this agreement.

I. CONDITIONS OF SUBMISSION

A. Scope of Agreement: Unless otherwise specified, the terms of this Agreement apply to the Manuscript and any and all supplemental digital content (SDC) (audio, video, and other data files, the forms of which may vary) submitted by the author(s) for online publication with the Manuscript. The Manuscript and the SDC hereafter are referred to collectively as the "Work."

B. Sole Submission of Original Work: Each author represents and warrants that the Work is the author's original work (except for any material from previously copyrighted works obtained by permission or assignment and credited as such); that none of the information has been published previously or is under consideration by another publication; that upon publication, the Work shall not infringe any copyright, invade any right of privacy, contain any libelous material, or infringe or violate any other right of any person or entity. Each author represents and warrants that the following is a complete description of all material contained in the Work, including any SDC, which is not original and/or in which the authors are not able to transfer copyright (attach additional pages if necessary):

C. Approval Verification: Each author represents and warrants that investigations involving human or animal subjects received the appropriate institutional approval and agrees to provide evidence of such approval if requested. In addition, each author agrees to produce the data on which the Work is based for examination by the editors or their assignees should they request them.

D. Authorship Certification: Each author represents and warrants that he or she has fulfilled the definition of authorship given in the Instructions for Authors, participated in this study sufficiently to ensure its integrity and to take public responsibility for it, and approved the Work submitted to *Obstetrics & Gynecology*. Each author represents and warrants that all persons who contributed to the work reported in the Manuscript are identified as authors or are included in the acknowledgements.

E. Conflict of Interest Disclosure: Each author represents and warrants that he or she has no financial affiliation (eg, employment, direct payments, stock holdings, retainers, consultancies, patent-licensing arrangements, or honoraria) or involvement within the last 3 years with any commercial organization with potential financial interest in the subject or materials discussed in the Work, except as disclosed in an attachment to this document. Any other potential conflict of interest also is disclosed.

F. Journal Research: Works under consideration are sometimes enrolled in *Obstetrics & Gynecology* research projects. When this is done, the identity of individual Works and authors is kept confidential. Each author hereby give consent for the Work to be used in such research.

G. Compliance with Privacy Laws: Each author represents and warrants that the Work complies with all applicable patient privacy laws (such as the Health Insurance Portability and Accountability Act of 1996, the regulations promulgated thereunder, and the Health Information Technology for Economic and Clinical Health Act of 2009, if applicable) and that the authors have obtained, prior to submission, any necessary patient authorizations. The corresponding author shall maintain such authorizations and shall inform the journal in a timely manner of any revocation by a patient of such authorization.

H. Identification of Patients: Each author represents and warrants that he or she has obtained a signed consent form from each patient described in a case report and each patient who is identifiable in a photograph or video. *Obstetrics & Gynecology* reserves the right to review the signed forms. Should the editorial office request copies of the signed forms, the corresponding author shall submit the forms in a timely manner.

I. Acknowledgments: Each author represents and warrants that all persons who contributed to the work reported in the Manuscript, but not sufficiently to be authors, are named in the acknowledgments and have given written permission to be named.

J. Manuscript Preparation Assistance: Each author represents and warrants that any and all document preparation assistance, including but not limited to topic development, data collection, analysis, writing, or editorial assistance, is disclosed in the acknowledgments.

K. Consideration: Each author hereby acknowledges that the publication of the Manuscript in a forthcoming edition of *Obstetrics & Gynecology* constitutes the sole consideration due to each of the authors under this Agreement.

I. Indemnification and Hold Harmless: Each author hereby agrees to indemnify, defend, and hold harmless the American College of Obstetricians and Gynecologists (the College), its directors, officers, agents, employees, affiliates, successors and assigns from and against all claims, actions, suits or other proceedings, and any and all losses, judgments, settlements, damages, expenses, liabilities (including liability assumed in contractual agreements with third parties), or other costs (including reasonable counsel fees, disbursements, and costs of settlement), arising from or in any way relating to the Work, or any breach, or alleged breach by any author of any of the provisions of this Agreement. If the Authors fail to promptly or diligently pursue any defense of any indemnified party, the indemnified parties, or any of them, may assume such defense at the expense of the Authors.

II. COPYRIGHT

A. Copyright Transfer: Each author hereby assigns, transfers, and otherwise conveys all rights, title, and interest in the Work, including all copyright ownership worldwide, in all languages, and in all forms of media now or hereafter known, including electronic media such as CD-ROM, Internet, and Intranet, for the full term and all extensions and renewals thereof, to the College. Each author acknowledges that, in addition to publishing the Work in *Obstetrics & Gynecology*, the College may use, publish, and distribute the Work in any manner or media in its sole discretion including, but not limited to, licensing the SDC to YouTube for inclusion on *Obstetrics & Gynecology's* YouTube channel. If the College and the editor should decide for any reason not to publish the Work in *Obstetrics & Gynecology*, one of them shall give prompt notice of the decision to the corresponding author, and this Agreement and each of the rights and obligations described herein shall terminate.

B. Work Made For Hire: If this Work, or any portion of it, was written in the course of the author(s)' employment as a "Work Made for Hire" as defined under U.S. Copyright Law, the Work is owned by the company/employer. The author must check the "Work for Hire" box after his or her signature and an authorized representative of the company/employer must also sign this form and state his or her title. In such case, the company/employer hereby assigns to the College, during the full term of copyright, all copyright in and to the submitted Work for the full term of copyright throughout the world as specified in Section II, paragraph A above.

C. Government Employees: If any author created this Work, or a portion of it, in the course of that author's employment by the United States Government, that author must check the "Government" box after his or her signature. A work prepared by a government employee as part of his or her official duties is called a "work of the U.S. Government" as it is not subject to copyright and the provisions of this Section II, paragraphs A and B, do not apply. If it is not prepared as part of the employee's official duties, it may be subject to copyright.

III. DEPOSIT OF POSTPRINTS

A. Author(s) Posting of Articles to an Institutional Repository: *Obstetrics & Gynecology* will permit the author(s) to deposit a "postprint" (defined as the final Manuscript after peer-review and acceptance for publication, but prior to the publisher's copyediting, design, formatting, and other services) on his/her personal Web site, university's institutional repository, or employer's intranet, subject to the following:

1. The author(s) may only deposit the postprint, as designated by *Obstetrics & Gynecology*, and not any other version of the Manuscript.
2. The author(s) shall not update the postprint text or replace it with a proof or with the final published version.
3. Except as provided in this Section, the author(s) shall not include the postprint or any other version of the article in any commercial website or in any repository other than as specified above.
4. The author(s) shall not display the postprint until 12 months after publication of the final article.
5. The author(s) must attach the following notice to the postprint: "This is a non-final version of an article published in final form in (provide complete journal citation)."
6. The author(s) shall provide a link in the postprint to the *Obstetrics & Gynecology* Internet site.

B. Compliance With Certain Research Funding Agency Requirements:

1. Authors of articles based on research funded by the National Institutes of Health (NIH), the Howard Hughes Medical Institute (HHMI), or the Wellcome Trust, are subject to the following policies:
 - a. NIH requires authors to deposit postprints based on NIH-funded research in the National Library of Medicine's (NLM) PubMed Central (PMC) within 12 months after publication of the final article.
 - b. HHMI requires authors to deposit postprints based on HHMI-funded research in NLM's PMC within 6 months after publication of the final article.
 - c. The Wellcome Trust requires authors to deposit postprints based on research funded by the Wellcome Trust in UK PubMed Central within 6 months after publication of the final article.
2. If the author(s) indicate at the bottom of this Agreement that the Manuscript is subject to a public access requirement, LWW will transmit the postprint of the Manuscript to NLM's PMC. The author(s) agree to the following:
 - a. Upon NIH request, it remains the legal responsibility of the author(s) to confirm with NIH the provenance of the Manuscript for purposes of deposit.
 - b. The author(s) shall not deposit their articles themselves.
 - c. The author(s) shall not alter the postprint already transmitted to NLM's PMC.

C. "Public Access Policy" Funding Disclosure

Please disclose below if you have received funding for research on which your article is based from any of the following organizations:

- ☐ National Institutes of Health (NIH)
- ☐ Wellcome Trust
- ☐ Howard Hughes Medical Institute (HHMI)
- ☐ Other (please list):
- ☐ Other (please list):

Each author must complete and sign a copy of this form to indicate his/her acceptance of the terms of this Agreement. Please check the applicable boxes.

Author:

Check if applicable (see Section II)

- ☐ Work for Hire (see Section II.B). Representative of company/employer must sign below.

Company/employer representative:	<input type="checkbox"/> Signature	<input type="checkbox"/> Printed Name	<input type="checkbox"/> Date
	<input type="checkbox"/> Title	<input type="checkbox"/> Company/Employer Name	

- ☐ Government (see Section II.C)

Disclosure of Potential Conflicts of Interest

- ☐ Yes, I have disclosed potential conflicts of interest (attached) (see Section I.E)

☒ No, I have no potential conflicts of interest to disclose (see Section I.E)

Authorship

In January 2014, *Obstetrics & Gynecology* adopted the ICMJE recommendations for authorship. *Note that you must meet a minimum of four criteria to be an author.* Individuals who do not qualify for authorship based on the ICMJE recommendations will be listed in the Acknowledgments. Indicate your contributions to the Work by checking the applicable boxes:

- ☒ 1a) Substantial contributions to the ^{or} conception or design of the Work
- ☐ 1b) Substantial contributions to the acquisition, analysis, or interpretation of data for the Work ^{and}
- ☒ 2) Drafting the Work or revising it critically for important intellectual content
- ☒ 3) Final approval of the version to be published
- ☒ 4) Agreement to be accountable for all aspects of the Work in ensuring that questions related to the accuracy or integrity of any part of the Work are appropriately investigated and resolved

Author Signature

Printed Name

Date

Author Agreement

Corresponding author:

EWELENA ROGOZINSKA 58 TURNER STREET, E12AB LONDON

Name

Address

+4420 7882 5881

+4420 7882 6047

E.A.ROGOZINSKA@QMUL.AC.UK

Telephone number

FAX number

E-mail address

**Title of Manuscript: ACCURACY OF ON-SITE TESTS TO DETECT ASYMPTOMATIC BACTERIURIA

**Name(s) of author(s): EWELENA ROGOZINSKA ; IN PREGNANCY: A SYSTEMATIC REVIEW

SANDRA FORMIA, JAVIER 2AMORA, AND META-ANALYSIS

LUCIANO MIGIMMI, KHALID KHAN

A signature below certifies agreement with the corresponding author designation and acceptance of this agreement. If there is more than one author, each must sign this agreement.

I. CONDITIONS OF SUBMISSION

A. Scope of Agreement: Unless otherwise specified, the terms of this Agreement apply to the Manuscript and any and all supplemental digital content (SDC) (audio, video, and other data files, the forms of which may vary) submitted by the author(s) for online publication with the Manuscript. The Manuscript and the SDC hereinafter are referred to collectively as the "Work."

B. Sole Submission of Original Work: Each author represents and warrants that the Work is the author's original work (except for any material from previously copyrighted works obtained by permission or assignment and credited as such); that none of the information has been published previously or is under consideration by another publication; that upon publication, the Work shall not infringe any copyright, invade any right of privacy, contain any libelous material, or infringe or violate any other right of any person or entity. Each author represents and warrants that the following is a complete description of all material contained in the Work, including any SDC, which is not original and/or in which the authors are not able to transfer copyright (attach additional pages if necessary):

C. Approval Verification: Each author represents and warrants that investigations involving human or animal subjects received the appropriate institutional approval and agrees to provide evidence of such approval if requested. In addition, each author agrees to produce the data on which the Work is based for examination by the editors or their assignees should they request them.

D. Authorship Certification: Each author represents and warrants that he or she has fulfilled the definition of authorship given in the Instructions for Authors, participated in this study sufficiently to ensure its integrity and to take public responsibility for it, and approved the Work submitted to *Obstetrics & Gynecology*. Each author represents and warrants that all persons who contributed to the work reported in the Manuscript are identified as authors or are included in the acknowledgements.

E. Conflict of Interest Disclosure: Each author represents and warrants that he or she has no financial affiliation (eg, employment, direct payments, stock holdings, retainers, consultancies, patent-licensing arrangements, or honoraria) or involvement within the last 3 years with any commercial organization with potential financial interest in the subject or materials discussed in the Work, except as disclosed in an attachment to this document. Any other potential conflict of interest also is disclosed.

F. Journal Research: Works under consideration are sometimes enrolled in *Obstetrics & Gynecology* research projects. When this is done, the identity of individual Works and authors is kept confidential. Each author hereby give consent for the Work to be used in such research.

G. Compliance with Privacy Laws: Each author represents and warrants that the Work complies with all applicable patient privacy laws (such as the Health Insurance Portability and Accountability Act of 1996, the regulations promulgated thereunder, and the Health Information Technology for Economic and Clinical Health Act of 2009, if applicable) and that the authors have obtained, prior to submission, any necessary patient authorizations. The corresponding author shall maintain such authorizations and shall inform the journal in a timely manner of any revocation by a patient of such authorization.

H. Identification of Patients: Each author represents and warrants that he or she has obtained a signed consent form from each patient described in a case report and each patient who is identifiable in a photograph or video. *Obstetrics & Gynecology* reserves the right to review the signed forms. Should the editorial office request copies of the signed forms, the corresponding author shall submit the forms in a timely manner.

I. Acknowledgments: Each author represents and warrants that all persons who contributed to the work reported in the Manuscript, but not sufficiently to be authors, are named in the acknowledgments and have given written permission to be named.

J. Manuscript Preparation Assistance: Each author represents and warrants that any and all document preparation assistance, including but not limited to topic development, data collection, analysis, writing, or editorial assistance, is disclosed in the acknowledgments.

K. Consideration: Each author hereby acknowledges that the publication of the Manuscript in a forthcoming edition of *Obstetrics & Gynecology* constitutes the sole consideration due to each of the authors under this Agreement.

L. Indemnification and Hold Harmless: Each author hereby agrees to indemnify, defend, and hold harmless the American College of Obstetricians and Gynecologists (the College), its directors, officers, agents, employees, affiliates, successors and assigns from and against all claims, actions, suits or other proceedings, and any and all losses, judgments, settlements, damages, expenses, liabilities (including liability assumed in contractual agreements with third parties), or other costs (including reasonable counsel fees, disbursements, and costs of settlement), arising from or in any way relating to the Work, or any breach, or alleged breach by any author of any of the provisions of this Agreement. If the Authors fail to promptly or diligently pursue any defense of any indemnified party, the indemnified parties, or any of them, may assume such defense at the expense of the Authors.

II. COPYRIGHT

A. Copyright Transfer: Each author hereby assigns, transfers, and otherwise conveys all rights, title, and interest in the Work, including all copyright ownership worldwide, in all languages, and in all forms of media now or hereafter known, including electronic media such as CD-ROM, Internet, and Intranet, for the full term and all extensions and renewals thereof, to the College. Each author acknowledges that, in addition to publishing the Work in *Obstetrics & Gynecology*, the College may use, publish, and distribute the Work in any manner or media in its sole discretion including, but not limited to, licensing the SDC to YouTube for inclusion on *Obstetrics & Gynecology's* YouTube channel. If the College and the editor should decide for any reason not to publish the Work in *Obstetrics & Gynecology*, one of them shall give prompt notice of the decision to the corresponding author, and this Agreement and each of the rights and obligations described herein shall terminate.

B. Work Made For Hire: If this Work, or any portion of it, was written in the course of the author(s)' employment as a "Work Made for Hire" as defined under U.S. Copyright Law, the Work is owned by the company/employer. The author must check the "Work for Hire" box after his or her signature and an authorized representative of the company/employer must also sign this form and state his or her title. In such case, the company/employer hereby assigns to the College, during the full term of copyright, all copyright in and to the submitted Work for the full term of copyright throughout the world as specified in Section II, paragraph A above.

C. Government Employees: If any author created this Work, or a portion of it, in the course of that author's employment by the United States Government, that author must check the "Government" box after his or her signature. A work prepared by a government employee as part of his or her official duties is called a "work of the U.S. Government" as it is not subject to copyright and the provisions of this Section II, paragraphs A and B, do not apply. If it is not prepared as part of the employee's official duties, it may be subject to copyright.

III. DEPOSIT OF POSTPRINTS

A. Author(s) Posting of Articles to an Institutional Repository: *Obstetrics & Gynecology* will permit the author(s) to deposit a "postprint" (defined as the final Manuscript after peer-review and acceptance for publication but prior to the publisher's copyediting, design, formatting, and other services) on his/her personal Web site, university's institutional repository, or employer's intranet, subject to the following:

1. The author(s) may only deposit the postprint, as designated by *Obstetrics & Gynecology*, and not any other version of the Manuscript.
2. The author(s) shall not update the postprint text or replace it with a proof or with the final published version.
3. Except as provided in this Section, the author(s) shall not include the postprint or any other version of the article in any commercial website or in any repository other than as specified above.
4. The author(s) shall not display the postprint until 12 months after publication of the final article.
5. The author(s) must attach the following notice to the postprint: "This is a non-final version of an article published in final form in (provide complete journal citation)."
6. The author(s) shall provide a link in the postprint to the *Obstetrics & Gynecology* Internet site.

B. Compliance With Certain Research Funding Agency Requirements:

1. Authors of articles based on research funded by the National Institutes of Health (NIH), the Howard Hughes Medical Institute (HHMI), or the Wellcome Trust, are subject to the following policies:
 - a. NIH requires authors to deposit postprints based on NIH-funded research in the National Library of Medicine's (NLM) PubMed Central (PMC) within 12 months after publication of the final article.
 - b. HHMI requires authors to deposit postprints based on HHMI-funded research in NLM's PMC within 6 months after publication of the final article.
 - c. The Wellcome Trust requires authors to deposit postprints based on research funded by the Wellcome Trust in UK PubMed Central within 6 months after publication of the final article.
2. If the author(s) indicate at the bottom of this Agreement that the Manuscript is subject to a public access requirement, LWW will transmit the postprint of the Manuscript to NLM's PMC. The author(s) agree to the following:
 - a. Upon NIH request, it remains the legal responsibility of the author(s) to confirm with NIH the provenance of the Manuscript for purposes of deposit.
 - b. The author(s) shall not deposit their articles themselves.
 - c. The author(s) shall not alter the postprint already transmitted to NLM's PMC.

C. "Public Access Policy" Funding Disclosure

Please disclose below if you have received funding for research on which your article is based from any of the following organizations:

- ☐ National Institutes of Health (NIH)
- ☐ Wellcome Trust
- ☐ Howard Hughes Medical Institute (HHMI)
- ☒ Other (please list): **WORLD HEALTH ORGANIZATION**
- ☐ Other (please list):

Each author must complete and sign a copy of this form to indicate his/her acceptance of the terms of this Agreement. Please check the applicable boxes.

Author:

Check if applicable (see Section II)

- ☐ Work for Hire (see Section II.B). Representative of company/employer must sign below.

Company/employer representative:	{	Signature	Printed Name	Date
		Title	Company/Employer Name	

- ☐ Government (see Section II.C)


Disclosure of Potential Conflicts of Interest

- ☐ Yes, I have disclosed potential conflicts of interest (attached) (see Section I.E)
- ☒ No, I have no potential conflicts of interest to disclose (see Section I.E)

Authorship

In January 2014, *Obstetrics & Gynecology* adopted the ICMJE recommendations for authorship. *Note that you must meet a minimum of four criteria to be an author.* Individuals who do not qualify for authorship based on the ICMJE recommendations will be listed in the Acknowledgments. Indicate your contributions to the Work by checking the applicable boxes:

- ☒ 1a) Substantial contributions to the conception or design of the Work
- ☒ 1b) Substantial contributions to the acquisition, analysis, or interpretation of data for the Work
- ☒ 2) Drafting the Work or revising it critically for important intellectual content
- ☒ 3) Final approval of the version to be published
- ☒ 4) Agreement to be accountable for all aspects of the Work in ensuring that questions related to the accuracy or integrity of any part of the Work are appropriately investigated and resolved


	ESELINA ROGOZIN'SKA	21-APR-2016
Author Signature	Printed Name	Date

C. "Public Access Policy" Funding Disclosure

Please disclose below if you have received funding for research on which your article is based from any of the following organizations:

- ☐ National Institutes of Health (NIH)
- ☐ Wellcome Trust
- ☐ Howard Hughes Medical Institute (HHMI)
- ☐ Other (please list):
- ☐ Other (please list):

Each author must complete and sign a copy of this form to indicate his/her acceptance of the terms of this Agreement. Please check the applicable boxes.

Author:		
<i>Check if applicable (see Section II)</i>		
<input type="checkbox"/> Work for Hire (see Section II.B). Representative of company/employer must sign below.		
<i>Company/employer representative:</i>	Signature	Printed Name
	Title	Company/Employer Name
<input type="checkbox"/> Date		
<input type="checkbox"/> Government (see Section II.C)		
Disclosure of Potential Conflicts of Interest		
<input type="checkbox"/> Yes, I have disclosed potential conflicts of interest (attached) (see Section I.E)		
<input checked="" type="checkbox"/> No, I have no potential conflicts of interest to disclose (see Section I.E)		
Authorship		
<small>In January 2014, <i>Obstetrics & Gynecology</i> adopted the ICMJE recommendations for authorship. <i>Note that you must meet a minimum of four criteria to be an author.</i> Individuals who do not qualify for authorship based on the ICJME recommendations will be listed in the Acknowledgments. Indicate your contributions to the Work by checking the applicable boxes:</small>		
<input type="checkbox"/> 1a) Substantial contributions to the conception or design of the Work		
<small>or</small>		
<input checked="" type="checkbox"/> 1b) Substantial contributions to the acquisition, analysis, or interpretation of data for the Work		
<small>and</small>		
<input checked="" type="checkbox"/> 2) Drafting the Work or revising it critically for important intellectual content		
<input checked="" type="checkbox"/> 3) Final approval of the version to be published		
<input type="checkbox"/> 4) Agreement to be accountable for all aspects of the Work in ensuring that questions related to the accuracy or integrity of any part of the Work are appropriately investigated and resolved		
	Javier Zamora	21/04/2016
Author Signature	Printed Name	Date

C. "Public Access Policy" Funding Disclosure

Please disclose below if you have received funding for research on which your article is based from any of the following organizations:

- ☐ National Institutes of Health (NIH)
- ☐ Wellcome Trust
- ☐ Howard Hughes Medical Institute (HHMI)
- ☐ Other (please list):
- ☐ Other (please list):

Each author must complete and sign a copy of this form to indicate his/her acceptance of the terms of this Agreement. Please check the applicable boxes.

Author:

Check if applicable (see Section II)

- ☐ Work for Hire (see Section II.B). Representative of company/employer must sign below.

Company/employer representative:	{	Signature	Printed Name	Date
		Title	Company/Employer Name	

- ☐ Government (see Section II.C)

Disclosure of Potential Conflicts of Interest

- ☐ Yes, I have disclosed potential conflicts of interest (attached) (see Section I.E)
- ☒ No, I have no potential conflicts of interest to disclose (see Section I.E)

Authorship

In January 2014, *Obstetrics & Gynecology* adopted the ICMJE recommendations for authorship. *Note that you must meet a minimum of four criteria to be an author.* Individuals who do not qualify for authorship based on the ICMJE recommendations will be listed in the Acknowledgments. Indicate your contributions to the Work by checking the applicable boxes:

- ☒ 1a) Substantial contributions to the conception or design of the Work
- or*
- ☐ 1b) Substantial contributions to the acquisition, analysis, or interpretation of data for the Work
- and*
- ☒ 2) Drafting the Work or revising it critically for important intellectual content
- ☒ 3) Final approval of the version to be published
- ☒ 4) Agreement to be accountable for all aspects of the Work in ensuring that questions related to the accuracy or integrity of any part of the Work are appropriately investigated and resolved

Author Signature

Printed Name

Date

Authors must submit their manuscripts online via Editorial Manager (ong.editorialmanager.com). Please do not send duplicate copies of the manuscript to the editorial office.

A COMPLETED CHECKLIST AND SIGNED AUTHOR AGREEMENT FORM MUST ACCOMPANY EACH MANUSCRIPT SUBMITTED TO *OBSTETRICS & GYNECOLOGY*.

Full Instructions for Authors are provided in the January and July issues or online (ong.editorialmanager.com).

General

- ☐ Author agreement forms are submitted.
- ☒ Potential conflicts of interest, whether of a financial or other nature, are disclosed by checking the appropriate box on the author agreement form and listing the potential conflicts in an attachment to the author agreement form.

- ☒ Pages are numbered consecutively.
- ☒ Lines are numbered consecutively throughout the manuscript using the line numbering feature available in word processing software.

- ☒ The manuscript is double-spaced.
- ☒ Only standard abbreviations and acronyms are included.

- ☐ For all original research, informed consent and institutional review board approval is addressed in the Methods. For case reports, a signed consent form has been obtained from each person who can be identified in a written description, photograph, or video (and is retained with your records).

- ☒ Permission to reprint material from another source in print and electronic form has been obtained.

- ☐ Clinical trials are registered in a public trials registry at or before the onset of patient enrollment.

- ☐ For randomized controlled trials, a completed CONSORT checklist accompanies the manuscript.

- ☐ For studies of diagnostic accuracy, a completed STARD checklist accompanies the manuscript.

- ☒ For meta-analyses and systematic reviews of randomized controlled trials, a completed PRISMA checklist accompanies the manuscript.

- ☐ For meta-analyses and systematic reviews of observational studies, a description of how the MOOSE guidelines are followed is included in the cover letter.

- ☐ For observational studies, a completed STROBE checklist accompanies the manuscript.

Cover Letter

- ☒ A cover letter is submitted indicating the intent to submit to *Obstetrics & Gynecology*.
- ☐ Publication elsewhere (including electronic media) of any of the material in

the manuscript submitted, other than an abstract of not more than 300 words, is identified and a copy of the other publication is submitted to the editorial office.

- ☐ If a version of the manuscript has previously been submitted for publication to *Obstetrics & Gynecology*, comments from the peer reviewers and an indication of how the authors have responded to these comments are included.

Title Page

- ☒ The title, author name(s) and major degree(s), affiliation(s), and the source(s) of funding for the work or study are provided.

- ☒ The name, address, telephone number, and e-mail address of the corresponding author are given.

- ☒ The manuscript title is no longer than 100 characters (letters and spaces) and does not contain any abbreviations or acronyms.

- ☒ A short title (no more than 45 characters) is provided at the bottom of the page for use as a running foot.

- ☒ The only acknowledgments are of financial or other substantive assistance. All individuals named in the acknowledgments have given written permission to be named.

Précis

- ☒ On a separate page, a statement of no more than 25 words is provided for the table of contents.

- ☒ Abbreviations or acronyms and commercial names are not used in the précis.

Abstract

- ☒ For original research reports, case reports, review articles, and procedures and instruments, the abstract is limited to 300, 125, 300, and 200 words, respectively, and follows the appropriate structured abstract format. Abstracts for Current Commentary articles should be a single paragraph of no more than 250 words that states what was done, what was found, and what the findings mean.

- ☒ Footnotes, references, and commercial names are not used in the abstract.

References

- ☒ References correspond to the examples available at ong.editorialmanager.com.

- ☒ References are identified using superscript or on the line within parentheses, eg, Levitz (3).

- ☒ Each reference is cited in the text.

- ☒ References are numbered consecutively in the order in which they appear in the text.

- ☒ Unpublished data, personal communications, statistical programs, papers presented at meetings and symposia, abstracts, letters, and manuscripts submitted for publication, are not listed in the reference list. Information from such sources is cited, if necessary, in text within parentheses.

Tables

- ☒ Each table is on a separate sheet of paper with its title.

- ☒ Tables are numbered with Arabic numerals.

- ☒ Each table contains all necessary information in order that it may stand alone, independent of the text.

- ☒ No table contains data that could be included in the text in several sentences.

Figures

- ☒ Digital art guidelines and artwork checklist on the journal's web site (ong.editorialmanager.com) have been consulted.

- ☒ Each figure is cited in the text.

- ☐ Letters and identifying marks (eg, arrows) are clear and sharp, and the critical areas of radiographs and photomicrographs are identified.

- ☐ In all cases (photograph or video) in which a human image is shown (in part or whole), written consent must also be obtained.

- ☒ Explanatory material appears in the accompanying legend and not on the figure itself.

- ☒ Legends are listed on one page at the end of the manuscript.





PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	3
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	4
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	4
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	4
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	5
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	5
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Supplementary materials
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	5
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	5-6
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	5
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	6
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	6
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2) for each meta-analysis.	6



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	6
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	6
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	7
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	Table 1
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	-
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	Supplementary materials
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	8-9
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	7-8, Figure 2
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	9
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	10
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	10-11
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	11-12
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	1

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.